

## The Goosander *Mergus merganser* breeding population expansion and trend in north-western Italy

LUCIO BORDIGNON<sup>1</sup>, MONICA CARABELLA<sup>2</sup>, WALTER GUENZANI<sup>2</sup>, MARCO GUERRINI<sup>3</sup>,  
NUNZIO GRATTINI<sup>4</sup>, ROBERTO LARDELLI<sup>5,6</sup>, GABRIELE PIOTTI<sup>3</sup>, CARLO PISTONO<sup>7</sup>,  
FABIO SAPORETTI<sup>2\*</sup>, MAURIZIO SIGHELE<sup>8</sup>, ANDREA TOSI<sup>9</sup>, ENRICO VIGANÒ<sup>10</sup>,  
GILBERTO VOLCAN<sup>11,12</sup>

<sup>1</sup> Gruppo Smergo - Via Vioglio 16, 13834 Soprana (BI), Italy

<sup>2</sup> Gruppo Insubrico di Ornitologia, c/o Civico Museo Insubrico di Storia Naturale di Clivio e Induno Olona Via Manzoni 21, 21050 Clivio (VA), Italy

<sup>3</sup> G.R.A., Gruppo Ricerche Avifauna - Brescia, Italy

<sup>4</sup> SOM, Stazione Ornitologica Modenese "Il Pettazzurro" - Comune di Mirandola (MO), Italy

<sup>5</sup> Ficedula, Associazione per lo studio e la conservazione degli uccelli della Svizzera italiana - Via Campo Sportivo 11 6834 Morbio Inferiore, Switzerland

<sup>6</sup> Swiss Ornithological Institute - Seerose 1, 6204 Sempach, Switzerland

<sup>7</sup> GOL, Gruppo Ornitologico Lombardo - Piazzale Veronica Gambara 5, 20146 Milano, Italy

<sup>8</sup> Verona Birdwatching, Italy

<sup>9</sup> 28887 Omegna (VB), Italy

<sup>10</sup> Polizia provinciale Nucleo Faunistico - Corso Matteotti 3, 23900 Lecco, Italy

<sup>11</sup> MUSE, Museo delle Scienze di Trento - Corso del Lavoro e della Scienza 3, 38122 Trento, Italy

<sup>12</sup> EBN-Italia, 30035 Moena (TN), Italy

\* Corresponding author: saporetif@gmail.com

**Abstract** – After first breeding in Italy in 1996 the Goosander has rapidly increased its breeding range in northern Italy: we analyzed the progressive expansion of Goosander in the lakes and rivers in a wide belt covering three-quarters of the sub-alpine Italian area (~45°N Lat). The study period covers the nesting seasons from 2010 to 2017. Each water body was assigned to a local coordinator that oversaw the monitoring operation during the breeding season, through a standardized census realized from the shores of each potential breeding area in the first week of June. Overall we censused a total of 282 broods (or families) in the whole period, distributed over 8 different water bodies, with the bulk of the population concentrated in the largest lakes, Maggiore, Como and Garda. The first census in 2010 estimated 12 broods while the maximum of 51 broods was reached in the last year of the survey (2017) suggesting a still ongoing increasing population trend. The number of chicks per brood ranged between 1 and 19, with a mean value of  $6.9 \pm 0.22$ . In literature brood size above 14 chicks are considered as due to brood amalgamation, which may therefore occur in north-Italian lakes. From 2011 the number of families was almost stable on the west of the study area (Lake Maggiore), while a non-significant decrease was revealed in the Lake Como, and a steady increase in the eastern Lake Garda. Overall, our findings describe the occurrence of a successful and still ongoing colonization towards east, possibly harbinger of further expansion.

**Key-words:** breeding range, population trend, brood size, Italy.

### INTRODUCTION

The Goosander breeds in a wide area of central and northern Europe, ranging from latitude 50° to 70° N. Below such limit, in the 70s, the species was confined as a breeder to the Swiss lakes and to the Bavaria region in Germany (Cramp & Simmons, 1977) which host a biogeographically isolated population (Hefti-Gautschi *et al.*, 2008). Out of the main

breeding area scattered couples are found in France, Slovenia, Ukraina and, in Southern Europe, in Albania and Greece (Hagemeyer & Blair 1997). An expansion process is underway in South (Switzerland and Italy) and Eastern Europe, in the Balkans, from the Carpathian mountains as far as Greece. In the lakes of the Dinaric Alps, in Western Serbia, the first female with chicks was observed in 1987, while in Bosnia-Herzegovina the first brood was registered

in 2005 (Marinković, 2008). Similarly a rapid colonization has been observed in the rivers of the Carpathian Mountains, in several areas and different countries (Czech Republic, Poland, Hungary, Romania), summing up 240 to 360 breeding pairs (Kaitoch & Bobrek, 2014). In the southern Balkans a small breeding population was known to persist from the begin of the twentieth century (Cramp & Simmons 1977, Scott & Rose 1996): in the period 2011-2015 a new census was conducted in the area comprising Macedonia, Albania and Greece, in the Ohrid and Prespa Lakes, that revealed a population of 15-25 breeding pairs (Catsadorakis *et al.* 2016). From '70s onwards, till the beginning of the 21st century, the Alpine population of Bavaria and Switzerland began to grow (BirdLife 2004, Schmid *et al.* 1998). The Swiss population was estimated at 490-670 pairs in 1998 (Keller & Gremaud 2003), further increasing at 600-800 pairs in the 2013-2016 period, when the species spread in the southern and eastern lakes of the country (Keller in Knaus *et al.* in press). The first brood in the Swiss portion of the Lake Maggiore was observed in 2003 (Volet & Burkhardt 2004). In Italy, at the beginning of the twentieth century, the Goosander was considered as a rare vagrant (Giglioli 1889, Arrigoni degli Oddi 1929), appearing only during migration and in winter: the few data were limited to Lake Garda, along the central trait of the Po river and in the Venetian lagoon (Arrigoni degli Oddi 1929). The first pair was found nesting in Italy in 1996, in the small artificial Lake Corlo in the municipality of Belluno in the Veneto Region, north-eastern alps (Zenatello *et al.* 1997), followed by a new pair on the western side of Lake Maggiore in 1998 (Bordignon 1999). It should be stressed that the two nesting localities are ~ 200 km away, and that many potentially breeding sites, as lakes and rivers, are found in this area. In 2002 the species was found breeding further east in northern Italy, in the Friuli Venezia Giulia region, along the Isonzo river that runs along the Slovenian border (Felcher & Utmar 2004) and, from 2000 to 2008, in small lakes and rivers (Piave and Brenta) of the Alpine sector of the Veneto and the Friuli Venezia Giulia region (Zenatello *et al.* 2009). The Italian breeding cohort of the species, from the beginning of the present century, revealed a positive trend, colonizing other lakes and rivers in northern Italy: in 2003 the first broods were observed in the eastern (Gagliardi *et al.* 2007) and northern (Volet & Burkhardt 2003) sectors of Lake Maggiore, followed by the first brood in 2004 on Lake Iseo (Bordignon, Pirola & Viganò, *pers.comm.*) joined in 2005 by the first on Lake Como (Viganò *et al.* 2006). Subsequently, further east, nesting pairs were found in 2010 (Gargioni & Piotti 2013) on the western shore of Lake Garda (the largest lake in Italy), followed respectively in 2014 and 2015, on the northern and eastern shores of the

same lake (Sighele *et al.* in press), with scattered pairs in other lakes and rivers of the pre-Alpine area. The Goosander nests in holes utilizing both natural and artificial cavities, preferably located in cliffs and rock walls, though not excluding buildings, abandoned Black Woodpecker's (*Dryocopus martius*) or nest-boxes (Cramp & Simmons 1977). Scarcity of nesting sites may induce females to share the same nest with other females: a reproductive tactic that has been defined as pre-hatch brood amalgamation (Eadie *et al.* 1988) or Conspecific Nest Parasitism (CNP, Pöysä 2006); a CNP threshold value of 14 chicks has been proposed by Eriksson & Niittyla (1985). Similarly to other cavity-nesting ducks such as, for example, the Wood Duck (Jones & Leopold 1967) or the Goldeneye (Eriksson & Andersson 1982, Pöysä & Pöysä 2002): for the Goosander there is evidence of a nest with 39 eggs (Géroudet 1985). The progressive colonization of the species in northern Italy prompted the creation of a voluntary monitoring group in 2010, the "Goosander Group", formed by several ornithologists and birdwatchers in northern Italy, that has since then coordinated the census of the breeding population in the study area. Our main interest was to assess

- 1) the current geographic breeding range of the species,
- 2) the population expansion and trend in lakes and rivers in north-western Italy,
- 3) the brood size of the families.

## MATERIAL AND METHODS


### Study area

The survey was conducted from 2010 to 2017 over almost the north western Italy (~ 80,000 km<sup>2</sup>) and focusing on 16 lakes and 6 main rivers (Tab. 1) of the Piedmont, Lombardy, and partly of the Trentino Alto Adige and Veneto regions bordering on Lake Garda, stretching from the pre-Alpine mountains to the Po Valley, north of the river Po (Fig. 1). The westernmost tip corresponds to Lake Avigliana (n. 1, Lat: 45.053 N, Long: 7.390E) in the Piedmont region, whereas the easternmost extreme is located at the northern tip of lake Garda (n. 15, Lat: 45.867, Long: 10.876) in Trentino Alto Adige. The northernmost part is located at Lake Mezzola (n. 7, Lat: 46.212 N, Long: 9.445 E), just north of lake Como; the southern one at the Lakes of Mantova (n. 16, Lat: 45.146 N, Long: 10.809E).

The lakes range from very small and shallow, bordered with reed beds and woods, with low depth, to the largest ones in Italy: Orta, Maggiore, Lugano, Como, Iseo and Garda (with the latter having the largest surface: ~370 km<sup>2</sup>) which are characterized mainly by mountain rocky coasts, dotted with built up areas overlooking the lake bor-

**Table 1.** Lakes and rivers that were monitored in the period 2010-2017, ordered from west to east, with mean altitude, surface and length. The main tributaries were also monitored on Lake Maggiore. The asterisk (\*) indicates occupied sites.

Lakes			Rivers	
sites	mean altitude	surface (km <sup>2</sup> )	sites	length (km)
1 Avigliana	354	1,5	17 Dora Baltea	168
2 Candia	226	1,52	18 Sesia	140
3 Viverone	230	5,8	19 Ticino*	110
4 Orta*	290	18,2	20 Adda*	164
5 Maggiore ( <i>Verbano</i> )*	193,8	212	21 Oglio	95
6 Lugano ( <i>Ceresio</i> )	270	48,9	22 Mincio	75
7 Mezzola	200	5,9		
8 Como ( <i>Lario</i> )*	199	145		
9 Montorfano	397	0,46		
10 Alserio	260	1,23		
11 Pusiano	257	5,2		
12 Annone	224	5,71		
13 Garlate*	198	4,64		
14 Iseo ( <i>Sebino</i> )*	185	65,3		
15 Garda ( <i>Benaco</i> )*	65	370		
16 Lakes of Mantova	18	6,21		



**Figure 1.** Map of the study area in northern Italy, with the 16 lakes and 6 river numbered progressively from west to east.

ders, with only a small amount of reed beds/woods, generally at the southern tip.

### Methods

Our census protocol required a local coordinator for every water body: for the largest ones (Maggiore, Como and Garda), the western and eastern sides were further subdivided between a minimum of two or more local coordina-


tors, to cover the entire length of the suitable shores; in the case of Lake Verbano there was a further subdivision because the northern part is Swiss territory. In each water body the broods were actively searched every year from May to July: this time frame encompasses the main period of the appearance of the females with chicks (a type of group known as “family parties”), in accordance with the census work already carried out in nearby Switzer-

land (Keller & Gremaud 2003). Every year a simultaneous census of the family parties was conducted at the 22 areas in the first week of June, that was utilized to compare the progress of the colonization in the various water bodies. The shores of lakes and rivers were actively searched (from 8.00 a.m. to 13.00 p.m.) from vantage points with binoculars and telescopes by monitoring groups formed of a minimum of two persons; in some years, on the largest lakes, we also have been supported by local management authorities who made a patrol boat available to scan the shores. During the census the position of every female with chicks was mapped: we consider a brood or family (female with chicks) as those one formed by chicks of the same age, registering different families if the observed age was clearly detectable. The only exception to this protocol regards the Lake Maggiore, on the Lombardy side from 2015 to 2017 (when the patrol boat was no longer available), and the Swiss part from 2010 to 2017, where observations on families were made not only in one day as in the previous years but encompassing the months of May and July. The numeric occurrence data for the Swiss side were resumed from the Ornitho.ch files of the Swiss Ornithological Institute: to avoid the possibility of double counts on different days, for assessing the number of broods we considered only the first observation of a family in the specified 1x1 km cell of the grid, disregarding further observations in the same cell. Following this method, the number of observed families might be slightly underestimated, so that the overall census should be considered as a conservative estimation. The Italian distribution of the Goosander for the period 2005-2017 was derived from the online map of the Italian Atlas of Breeding Birds (UTM cartographic grid of 10x10 km) available on the web-portal “ornitho.it”. The Goosander has synchronous hatching and it has been established that broods may be gathered where numerous so that excessively large broods may originate from more than one female (Cramp & Simmons 1977) and we adopted the threshold value of 14 chicks as proxy for brood amalgamation (Savard 1987, Eadie *et al.* 1988). Every year, at the end of the monitoring session, the counts were transmitted to the project coordinator (Lucio Bordignon) for data storage. Statistical analyses and graphs were made with R software (R Development Core Team 2018, version 3.4.4) using “Stats”, “lme4”, “sjstats” and “ggplot2” packages (Tibco Enterprise 2017, Bates *et al.* 2018, Lüdecke 2018, Wickham & Chang 2016). We used a linear regression model to test the increase of family parties in the years for the whole study area and for the four largest water bodies (lakes Maggiore, Como, Iseo and Garda), displayed with a Kernel Regression Smoother. After fitting the model, we checked the distribution of the

standardized residual with a Normal Q-Q plot. To assess whether brood size varied across years, we fitted a Generalized Linear Mixed Model (package “lme4”) with number of juveniles per brood as the dependent variable, year (entered after standardization) as the predictor, and site as a random factor, to correct for spatially non-independent records. Given that the use of a Poisson error led to an overdispersed model (with residual deviance much higher than residual degrees of freedom, and an overdispersion level significant at  $P < 0.001$  according to the ‘overdisp’ command in the ‘sjstats’ package), we fitted a negative binomial model, which was not overdispersed ( $P = 0.357$  according to the same test). Then, we assessed the significance of the year effect and compared the value of the Akaike’s Information Criterion (AIC) of the model with and without year, respectively.

## RESULTS

Overall in eight years of census we found a total of 282 broods distributed in eight water bodies, representing the 36.4% of the monitored sites. The yearly number of family parties showed a steady increase during the study period (Fig. 2) statistically significant (linear model  $R^2 = 0.807$ ;  $\beta = 11.66 \pm 2.32$ ;  $t = 5.02$ ,  $P = 0.002$ ); beginning with 12 families in 2010 and reaching a maximum of 51 in 2017 (Tab. 2). Only the four major lakes (Maggiore, Como, Iseo and Garda) were occupied all the years; in 2013 families were observed for the first time on two rivers (Ticino and



**Figure 2.** Regression line for the total number of families per year.

**Table 2.** Counted number of families (female with chicks) per year.

year	2010	2011	2012	2013	2014	2015	2016	2017
n. families	12	28	28	31	49	41	42	51

Adda, respectively sites n. 19 and 20) and, from 2015 till 2017, on the river Toce that belongs to the water catchment of Lake Maggiore. In 2016 a territorial pair was observed along the Sesia river (n. 18) and again two pairs in 2017, but no evidence of breeding was discovered in both years. In 2017 one family party was observed respectively on lakes Orta (site n. 4) and another one in Garlate (site n. 13): in the first site already in 2015 a pair was observed without confirming the breeding. Lake Garlate is a small lake, at the southern tip of Como lake, with heavily built-up shores. From 2010 to 2014 the bulk of the population (84.3%) was concentrated in two lakes (Tab. 3), Maggiore (site n. 5) e Como (site n. 8). Lake Iseo (site n. 14) has always hosted a low number of families during the study period, with a maximum of 3 families in 2016, whereas Lake Garda (site n. 15) showed a positive trend (Fig. 3), starting with the first observed brood in 2010, and reaching the number of 16 families in 2017 (linear model  $R^2 = 0.83$ ;  $\beta = 4.87 \pm 0.90$ ;  $t = 5.42$ ,  $P = 0.002$ ); The other three sites (n. 5, 8, 14) do not reveal a statistically significant trend, with fluctuating number of broods, more pronounced on Lake Como in the last two years of the census. The number of chicks per brood ranged from 1 to 19, with a mean value


of 6.94 (ES = 0.22; median = 6, N= 282 broods), but the frequency distribution resulted right-skewed so that small broods appeared more abundant (Fig. 4). In the 78.4% of the cases (N = 211) , broods comprised up to 9 chicks, while the 21.6% comprised 10 or more chicks. Only 15 broods (5.31% of the total) showed a number of chicks equal or higher than the threshold value stated for Con-specific Nest Parasitism ( $\geq 14$  chicks); the median value of chicks per broods decreased during the first three years stabilizing from 2013 onwards. To analyze temporal variation in brood size we fitted a negative binomial model with year as predictor: the year effect was not significant ( $\beta \pm ES: -0.01 \pm 0.02$ ;  $P = 0.579$ ) and the model with year had a higher AIC ( $\Delta AIC=1.87$ ) then the null model.

## DISCUSSION

The European population of the Goosander is estimated at 66,800-103,000 pairs, with the stronghold of the specie being hosted in Finland (20,000-30,000 pairs or 30% of the European population) and Sweden (27,000-42,000 pairs or 41% of the European population). The Goosander is not a threatened species and the regional assessment, in the frame of the IUCN Red List Status, defines the species as Least Concern (LC, BirdLife International 2015). In the long (1980-2012) and short-term (2001-2012), the trend of the EU population is defined as “decreasing”, mainly on account of the negative trend of the Estonian and Finland population, with a magnitude that ranged from 31 to 49%.

**Table 3.** Total number of broods counted in the four major lakes, mean (ES) and median.  $R^2$  value for the linear model for the 8 years trend, with P value.


	Maggiore (site n. 5)	Como (site n. 8)	Iseo (site n. 14)	Garda (site n. 15)
2010	6	4	1	1
2011	14	12	1	1
2012	9	15	2	2
2013	10	14	2	2
2014	22	21	2	4
2015	11	19	2	7
2016	13	15	3	10
2017	14	15	1	16
<b>total</b>	<b>99</b>	<b>115</b>	<b>14</b>	<b>43</b>
<b>mean</b>	<b>12.4</b>	<b>14.4</b>	<b>1.6</b>	<b>5</b>
<b>ES</b>	<b><math>\pm 1.68</math></b>	<b><math>\pm 1.79</math></b>	<b><math>\pm 0.25</math></b>	<b><math>\pm 1.89</math></b>
<b>median</b>	<b>12</b>	<b>15</b>	<b>2</b>	<b>3</b>
$R^2$ - Linear model	0.18	0.43	0.17	0.83
P value	0.295	0.077	0.31	0.002



**Figure 3.** Smoothed trend-lines of the number of families on lakes Maggiore, Como and Garda.

In this context emphasis must be placed on the assessed positive short-term trend of the majority of the central and eastern European countries, that is similar, but smaller in size, to that observed in the southern countries (BirdLife International 2015). Concerning the relatively recent colonization of Italy Gustin *et al.* (2016) did not provide a Favourable Reference Value for the Goosander in Italy. The European guidelines aimed at evaluating and monitoring the conservation status of the species, require that all the Member States, as a long-term objective that takes into consideration the population size, range and habitat, provide a “Value” that may represent a favourable status of species. This “Value” may be translated as numbers of pairs per unit area for those specie regularly breeding in Italy and Gustin *et al.* (2016) indicates a Minimum Viable

Population of 300 pairs as a long-term conservation objective. The recent Italian known breeding range of the species, as results from the map available from the ornitho.it data-base ([www.ornitho.it](http://www.ornitho.it); accessed on 28 February 2018, for the period 2005-2017), regards only the northern part of the country, from River Toce and Lake Orta on the west (site n. 4), up to the Slovenian border on the east, with observations regarding 88 10x10 km atlas square, of which 55 are related to breeding individuals. Only in 2017, in our study area we had a minimum of 51 females with chicks, corresponding to 21 10x10 atlas square with successful nesting, matched in north-east Italy (outside our study area) by 9 atlas square with a minimum of 10 families: hence we obtain a minimum of 60 nesting pairs. From these data we can estimate a minimum of 80-150 nesting pairs for Northern Italy, a three to five-fold increase of the previous published reference that assessed 22-29 pairs for the period 2008-2012 (Nardelli *et al.* 2015). The results of our survey reveal that the bulk of the Goosander’s population is still concentrated in the three largest lakes (Maggiore, Como and Garda), with smaller lakes and rivers being colonized by a low number of pairs. Since 2014 onwards the breeding records on Lake Garda rose to the same level of Lakes Maggiore and Como, but the spreading of the nesting pairs on Lake Garda appears to be more rapid than the first two. From 2015 onward the breeding population in the western lakes (Maggiore and Como) appears to be stable: in the first lake the Goosander is becoming increasingly frequent in the southern part of the water body, that presents different characteristics to those of the northern sector, where the rocky shore is predominant. In the southern part, partly urbanized, the species is also more frequent (Saporetto 2018) in the reeds areas of the Special Protection Area “IT2010502 Canneti del Lago Maggiore”, a marshland-woodland site, where the first brood was observed in 2017. Other broods were observed, from 2015 to 2017, both in-



**Figure 4.** Frequency distribution of the number of chicks per broods with mean (indicated by dashed line).

**Table 4.** Values of mean brood size available in literature and during the present study.

paper	country	mean brood size	ES	number of families
Bauer & Zintl (1974)	Germany (Bayern)	7		44
this paper	Italy	6.9	± 0.99	282
Catsadorakis <i>et al.</i> (2016)	Albania F.Y.R. of Macedonia Greece	6.4	± 0.56	38
Keller & Gremaud (2003)	Switzerland	6.3 6.5	± 0.40 ± 0.41	min. 241 max. 255
Marinkovic <i>et al.</i> (2008)	Serbia Bosnia-Erzegovina	6.5		126
Zenatello <i>et al.</i> (2009)	Italy	7.4	± 3.02	6

side and outside our study area, demonstrating the continuous colonization process that involves smaller rivers. The mean number of chicks per brood observed in the present survey is 6.9 (ES ± 0.99) and this value is similar with those available from literature, whose values ranging from 6.3 to 7.4 (Tab. 4). The expansion of the breeding range is matched by the trend of the wintering individuals as obtained by IWC census data, despite a relatively recent discovery of a north-east range shift in wintering individuals (Musilová *et al.* 2009, Lehtikoinen *et al.* 2013) connected with climate change. On this aspect we may assume a different wintering behaviour between the Alpine population and the north-European population. Indeed, the wintering numbers in the Lombardy region, passed from 11 individuals in 2002 (Vigorita *et al.* 2002) to a maximum of 371 in the year 2016 (Longoni & Fasola 2016), taking into account a period of 15 years.

**Acknowledgements** – This survey was supported by all those who participated over the years. We are grateful to: Davide Aldi, Gianfranco Alessandria, Roberto Aletti, Roberta Allievi, Sergio Baratelli, Gabriele Baruffaldi, Stefano Bellintani, Stefano Benini, Luca e Vittorio Bergamaschi, Angelica e Marco Bergomi, Eros Bernardara, Martino Bertalla, Paolo Bertini, Federica Bertola, Roberto Bertoli, Cesare Bertoni, Radames Bionda, Roberto Boscai ni, Anna Emma Botta, Gianni Bragante, Sonia Braghiroli, Roberto Brembilla, Paola Bressan, Massimo Brigo, Fabrizio Burlone, Martina Cadin, Ivan Callovi, Isabella Cappelletti-Arnold, Monica Carabella, Augusta Castiglioni, Daniela Casola, Piero Castelli, Nicola Chiavone, Silvio Colaone, Lorenzo Colombo, Andrea Comalini, Giancarlo Comincioli, Gianluca Cordioli, Dario Crinari, Federico De Pascalis, Francesca Diana, Markus Ehrenguber, Luca Faccioli, Paolo Faifer, Massimiliano Favoni, Nicola Ferrari, Giovanni Fontana, Emanuele Forlani, Diego Franchini, Alessandro Franzoi, Marco Fredi, Arturo Gargioni, Paolo Gialdi, Abramo Giusto, Nunzio Grattini, Walter Guenzani, Marco Guerrini, Claudio Isotta, Erika Kessler, Hannes Klopfenstein, Donatella Lampugnani, Roberto Lardelli, Franco Lorenzini, Aurelio Lubini, Fausto Lucani, Giuseppe Luoni, Roberto Lupi, Alessandro Madella, Stella Magrone, Susanna Malfer, Gianni Marcolli, Celestino Marelli, Julien Mazenauer, Stefania Mazzaracca, Margherita Mazzetti, Luca Mazzola, Paolo Merga, Alessandro Micheli, Annunziata

Monte, Thea Moretti, Massimo Mormile, Ettore Mozzetti, Marta Musatti, Giovanni e Maria Grazia Natale, Ostinelli-Cattin Orlando, Alison e Colin Parnell, Massimiliano Pasini, Angelo Pasqua, Maurizio Pasquali, Giuliano Pasquariello, Paolo Pedrini, Giacomo Pellegrini, Alice Pellegrino, Vincenzo Perin, Marina Pestoni, Federico Pianezza, Riccardo Pianezza, Gabriele Piotti, Stefano Pirola, Giuliana Pirotta, Pietro Pisano, Carlo Pistono, Dario Quaranta, Marco Ranaglia, Giovanni Ratti, Alvinio Ravasi, Giuseppe Redaelli, Nicola Redaelli, Marco Ricci, Paola Ricceri, Ettore Rigamonti, Ruggero Rizzi, Antonio Ronconi, Roberto Santinelli, Fabio Saporetti, Fabiano Sartirana, Chiara Scandolara, Maurizio Sighele, Nadir Signori, Giuseppe Speranza, Raffaele Spocci, Mauro Stangalini, Gerri Stefani, Paolo Stephani, Michael Straubhaar, Morena Stornetta, Karol Tabarelli de Fatis, Pietro Teichert, Michele Tognina, Mirko Tomasi, Damiano Torriani. Andrea Tosi, Damiano Torriani, Paolo Trotti, Giuliana Turconi, Lucio Uber, Michel-Thomas Usteri, Daniele Vezzoli, Andrea Vidolini, Andrea Viganò, Enrico Viganò, Gilberto Volcan, Fulvio Zanardini, Tonino Zarbo. We apologize to anyone whose name we may have unwittingly omitted from this acknowledgement. We thank Michele Panuccio and two anonymous referees for their constructive criticisms of the earlier draft. We are extremely grateful to Mattia Brambilla for his assistance with statistical modelling. We wish to thank the Swiss Ornithological Institute for permission to consult their data, particularly Verena Keller for her helpful comments relating to the Swiss data and for giving us permission to quote data from the new Swiss Atlas of Breeding Birds (in press); Chiara Scandolara for her supportive help and advice. MUSE (Museo delle Scienze, Trento) has promoted the fieldwork in the Trentino side of lake Garda. Our thanks are due to the wildlife wardens of Brescia, Como, Lecco, Sondrio and Varese for their invaluable and friendly cooperation. We are grateful to the following groups for their organizational support: Novara Bird-Watching, EBN Italia, GIO (Gruppo Insubrico di Ornitologia), GPSO (Gruppo Piemontese Studi Ornitologici “Bonelli”), GRA (Gruppo Ricerche Avifauna), GRAM (Gruppo di Ricerche Avifauna Mantovano). Many thank to Paul Nicholls for revising the English text.

## REFERENCES

- Arrigoni degli Oddi E., 1929. Ornitologia Italiana. U. Hoepli ed., Milano.
- Bates D., Maechler M., Bolker B., Walker S., Package “lme4”, 2018. Linear Mixed\_Effects Models using “Eigen” and S4. Vers. 1.1-17.
- Bauer U. & Zintl H., 1974. Brutvorkommen und Brutbiologie

- des Gänsesäger *Mergus merganser* in Bayern. Anz. Ornithol. Ges. Bayern 13: 71–86.
- BirdLife International, 2004. Birds in Europe: population estimates, trends and conservation status. BirdLife International Conservation Series, Cambridge, UK.
- BirdLife International, 2015. European Red List of Birds. Luxembourg: Office for Official Publications of the European Communities.
- Bordignon L., 1999. Prima nidificazione dello smergo maggiore in Piemonte. Riv. ital. Orn. 69: 218–220.
- Catsadorakis G., Avramoski O., Bojadzi A. & Nikolaou H., 2016. The status of an isolated population of goosander *Mergus merganser* in the Balkans. Wildfowl 66: 159–175.
- Cramp S. & Simmons K.E.L. (eds.), 1977. The birds of the Western Palearctic. Vol. 1. Oxford University Press, Oxford.
- Eriksson M.O.G. & Andersson M., 1982. Nest parasitism and hatching success in a population of goldeneyes *Bucephala clangula*. Bird Study 29: 49–54.
- Eriksson K. & Niittylä J., 1985. Breeding performance of the Goosander *Mergus merganser* in the archipelago of the Gulf of Finland. Ornith. Fenn. 62: 153–157.
- Eadie J.M., Kehoe F.P. & Nudds T.D., 1988. Pre-hatch and post-hatch brood amalgamation in North American Anatidae: a review of hypotheses. Canadian J. Zool. 66 (8): 1709–1721.
- Felcher L. & Utmar P., 2004. La nidificazione dello Smergo maggiore, *Mergus merganser*, nel Friuli-Venezia Giulia. Riv. Ital. Orn. 74: 69–71.
- Gagliardi A., Guenzani W., Preatoni D.G., Saporetto F. & Tosi G. (eds.), 2007. Atlante Ornitologico Georeferenziato della provincia di Varese - uccelli nidificanti 2003-2005. Provincia di Varese, Museo Insubrico St. Nat. Induno Olona, Univ. dell'Insubria di Varese, 295 pp.
- Gargioni A. & Piotti G., 2013. Prima nidificazione di smergo maggiore in provincia di Brescia. Natura Bresciana 38: 133–134.
- Géroudet P., 1985. Essai de synthèse sur l'évolution du Harle bièvre, *Mergus merganser*, dans le bassin du Léman. Nos Oiseaux 38: 1–18.
- Giglioli E.H., 1889. Avifauna Italica. Elenco sistematico delle specie di uccelli stazionarie o di passaggio in Italia. Le Monnier, Firenze.
- Gustin M., Brambilla M. & Celada C., 2016. Stato di conservazione e valore di riferimento favorevole per le popolazioni di uccelli nidificanti in Italia. Riv. ital. Orn. 86 (2): 3–58.
- Hagemeijer E.J.M. & Blair M.J. (eds.), 1977. The EBCC Atlas of European Breeding Birds: their distribution and abundance. T. & A.D. Poyser, London.
- Hefti-Gautschi B., Pfunder M., Jenni L., Keller V. & Ellegren H., 2008. Identification of conservation units in the European *Mergus merganser* based on nuclear and mitochondrial DNA markers. Conservation Genetics 10: 87–99.
- Jones R.R. & Leopold A.S., 1967. Nesting interference in a dense population of Wood Ducks. J. Wildl. Manage. 31: 221–229.
- Kajtoch Ł. & Bobrek R., 2014. Extension of Goosander *Mergus merganser* distribution into the Carpathian range. Wildfowl 64: 91–101.
- Keller V. & Gremaud J., 2003. Der Brutbestand des Gänsesägers in der Schweiz 1998. Orn. Beob. 100: 227–246.
- Knaus P., Antoniazza S., Wechsler S., Guélat J., Kéry M., Strelben N. & Sattler T., in press. Atlante degli uccelli nidificanti in Svizzera 2013-2016. Distribuzione ed evoluzione degli effettivi degli uccelli in Svizzera e nel Liechtenstein. Stazione Ornitologica Svizzera, Sempach.
- Lehikoinen A., Jaatinen K., Vähätalo A.V., Clausen P., Crowe O., Deceuninck B., Hearn R., Holt C.A., Hormman M., Keller V., Nilsson L., Langedoen T., Tománková I., Wahl J. & Fox. A.D., 2013. Rapid climate driven shifts in wintering distributions of three common waterbird species. Global Change Biology 19: 2071–2081.
- Longoni V. & Fasola M. 2016. Censimento Annuale degli Uccelli Acquatici Svernanti in Lombardia. Resoconto 2016. Regione Lombardia, Milano.
- Lüdecke D., Package “sjstats”, 2018. Collection of Convenient Functions for Common Computations. Vers. 0.15.0.
- Marinković S.P., Skorić S.B., Popović Z.S. & Nikčević M.V., 2008. Research on long-term colonization of Goosander (*Mergus merganser* Linnaeus, 1758) with reference to habitat availability. Arch. Biol. Sci., Belgrade 60 (3): 501–506.
- Musilová Z., Musil P., Poláková S. & Fuchs R., 2009. Wintering ducks in the Czech Republic: changes in their population trends and distribution. Wildfowl Special Issue 2: 74–85.
- Nardelli R., Andreotti A., Bianchi E., Brambilla M., Brecciaroli B., Celada C., Dupré E., Gustin M., Longoni V., Pirrello S., Spina F., Volponi S. & Serra L., 2015. Rapporto sull'applicazione della Direttiva 147/2009/CE in Italia: dimensione, distribuzione e trend delle popolazioni di uccelli (2008-2012). ISPRA, Serie Rapporti, 219/2015.
- Pöysä H. & Pöysä S., 2002. Nest-site limitation and density dependence of reproductive output in the common goldeneye *Bucephala clangula*: implications for the management of cavity-nesting birds. J. appl. Ecol. 39: 502–510.
- Pöysä H., 2006. Public information and conspecific nest parasitism in goldeneyes: targeting safe nests by parasites. Behav. Ecol. 17 (3): 459–465.
- R Development Core Team (3.4.4 2018). The R Foundation for Statistical Computing.
- Saporetto F., 2018. L'avifauna acquatica della ZPS IT2010502 “Canneti del Lago Maggiore”. Risultati del monitoraggio 2015-2017. Relazione tecnica non pubblicata.
- Savard J.P.L., 1987. Causes and functions of brood amalgamation in Barrow's Goldeneye and Bufflehead. Canadian J. Zool. 65 (6): 1548–1553.
- Schmid H., Luder R., Naef-Daenzer B., Graf R. & Zbinden N., 1998. Schweizer Brutvogelatlas. Verbreitung der Brutvögel in der Schweiz und im Fürstentum Liechtenstein 1993-1996. Schweizerische Vogelwarte, Sempach.
- Scott D.A. & Rose P.M., 1996. Atlas of Anatidae populations in Africa and Western Eurasia. Wetlands International Publication n. 41, Wetlands International, Wageningen, The Netherlands.
- Sighele M., Boscaini R. & Speranza G., in press. Prime nidificazioni di smergo maggiore, *Mergus merganser*, sulla sponda veronese del lago di Garda (Aves: Anatidae). Atti VIII Conv. Faunisti Veneti.
- Tibco Enterprise Runtime for R stats package. 2017. Vers: 4.3.0-44.
- Viganò E., Ornaghi F. & Pasquariello G., 2006. Prima nidificazione di Smergo maggiore *Mergus merganser* in Lombardia. Picus 32: 115–116.
- Vigorita V., Rubolini D., Cucè L. & Fasola M., 2002. Censimento annuale degli uccelli acquatici svernanti in Lombardia. Resoconto 2002. Regione Lombardia.
- Volet B. & Burkhardt M., 2003. Übersicht über das Brutgeschehen und andere ornitologische Ereignisse 2002 in der Schweiz. Orn. Beob. 101: 281–294.
- Wickham H. & Chang W., Package “ggplot2”, 2016. Create Elegant Data Visualizations Using the Grammar of Graphics. Vers. 2.2.1.
- Zenatello M., Basso S., Rasi S. & Tormen G., 1997. Prima nidificazione di Smergo maggiore in Italia. Riv. ital. Orn., 66 (1996): 207–210.
- Zenatello M., Bordignon L., Ventolini N., Utmar P. & Viganò E., 2009. Lo Smergo maggiore nidificante in Italia: 1996-2008. Alula 16: 491–496.